

Brake Herbicide for Cotton

Brake®
Herbicide

Broad-spectrum preemergence cotton herbicide

Provides residual control of annual grass and broadleaf weeds

- Excellent Pigweed (Palmer amaranth) control
- Annual grass control including resistant goosegrass and Italian ryegrass
- Great residual control in wet conditions
- Safest preemergence herbicide available in cotton
- A unique HRAC/WSSA Group 12 mode of action
- Compatible with other herbicides and liquid fertilizer
- Crop rotation flexibility for your farming operation



Brake® Herbicide Program

Start weed free, stay weed free. Use multiple modes of action. Overlap residuals.

1. Apply a preplant burndown to control existing weeds
2. Tank-mix 16 fl. oz. Brake® with another residual herbicide (e.g. Cotoran® or Reflex®) and a burndown (e.g. Gramoxone®)

Proven tank-mixes

- 16 fl. oz. Brake® + Cotoran® + Burndown
- 16 fl. oz. Brake® + Reflex® + Burndown
- 16 fl. oz. Brake® + Diruon® + Burndown
- 16 fl. oz. Brake® + Warrant® + Burndown

3. Apply Brake tank-mix at planting or within 36 hours of planting
4. Make a postemergence application within 18 days after planting containing an overlapping residual (e.g. Dual Magnum®) + non-residual(s) (e.g. Liberty®) compatible with your cotton variety
5. Scout and utilize additional postemergence applications with overlapping residuals as needed

Stop Weeds. Protect Cotton.

Plan Your Program, Work Your Program!

Crop Rotation Restrictions

Crops to be Planted	Minimum Rotation Interval (months after last Brake application) ¹		
	16 fluid ounces or less Brake [®] per acre	17 - 21 fluid ounces or less Brake [®] per acre	22 - 32 fluid ounces or less Brake [®] per acre
Cotton	0	0	0
Peanut	0	8	8
Rice	0	8	12
Wheat, barley and rye	8 ²	8	12
Sweet Potato	2	2	2
Soybean	2	4	12
Corn and sorghum	10 ³	12 ³	18
Alfalfa, sunflower and crops in the Solanaceae family (e.g. peppers, tobacco, tomatoes)	18	18	18

¹ West TX only add 4 months to rotational restrictions for all crops except cotton and peanut.

² If used as a cover crop (i.e. will not be harvested), wheat, barley, and rye may be planted 5 months after the last Brake application if no more than 16 fluid ounces were applied.

³ Minor chlorosis and stunting of corn and sorghum may occur, especially if heavy rainfall occurs during or soon after emergence, but plants typically outgrow these effects and develop normally. If this product is used for two consecutive years in the same field at a rate greater than 16 fl oz per acre, only plant cotton, peanut, rice, or soybean the following year in those fields.

Learn More



SePRO Ag, LLC Carmel, IN 46032 ag.sepro.com (844) 766 2727

SePRO Ag[™]
Upgrade Your Crop[™]

Always read and follow label directions. Brake is a registered trademark of SePRO Corporation. Diruon is a registered trademark of Loveland Products, Inc. Gramoxone, Dual Magnum, and Reflex are trademarks of a Syngenta Group Company. Cotoran is a registered trademark of an ADAMA Group Company. Liberty is a registered trademark of BASF. Warrant is a registered trademark of Monsanto Technology LLC. ©2024 SePRO Ag, LLC. New 03/14/2024.