

Supplemental Labeling



SePRO Ag, LLC

11550 N. Meridian Street, Suite 600

Carmel, IN 46032 USA

Brake[®]

EPA Reg. No. 67690-78-101232

Residual herbicide for use in rice to control a range of annual grass and small seeded broadleaf weeds including pigweed.

This label expires January 25, 2026 and must not be used or distributed after this date.

KEEP OUT OF REACH OF CHILDREN CAUTION / PRECAUCIÓN

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Brake[®] before applying. Carefully follow Precautionary Information and applicable use directions.
- Use of Brake according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Brake.

The use directions are based on the results of product testing programs across a wide variety of environments and varieties/cultivars. However, not all varieties/cultivars have been tested across all environments, including potential tank-mix combinations and sequential pesticide applications. Consult your local SePRO Ag, LLC representative to determine if information exists for the particular use (e.g. variety/cultivar, crop age, tank-mixes, etc.), especially if the environment contains coarse textured soils with a soil pH greater than 7.5. If no information is available, the user assumes all risk and liability associated with commercial applications if the particular use (e.g. crop variety/cultivar, crop growth stage, tank-mixes, etc) is not first tested on a small area under local conditions.

Rice Use Restrictions

- **DO NOT** apply by air.
- Chemigation: **DO NOT** apply through any type of irrigation system.
- **DO NOT** apply more than 16 fluid ounces of this product per acre per application (equivalent to 0.15 lb fluridone per acre) in rice.
- **DO NOT** apply more than 16 fluid ounces of this product per acre per year (equivalent to 0.15 lb fluridone per acre) in rice.
- **DO NOT** make postemergence applications prior to the 3-leaf stage.
- **DO NOT** apply to water-seeded rice.
- **DO NOT** apply to ratoon rice.
- **DO NOT** apply to wild rice.

- **DO NOT** apply where runoff or irrigation water may flow onto agricultural land other than cotton or rice fields.
- **DO NOT** use treated water for any form of irrigation.
- **DO NOT** use treated water for hydroponic farming.
- **DO NOT** apply to salt/brackish water.
- **DO NOT** apply this product as a standalone residual herbicide to the same field more than 2 years in a row.
- **Preharvest Interval:** Do not apply within 30 days of harvest.

Rice Use Precautions

- Application of Brake followed by conditions that do not promote normal crop growth and development, or cause stress (e.g. cold temperatures), may result in crop injury.
- Before applying to rice, verify with your local SePRO Ag, LLC representative the selectivity of this product on each variety that will be planted to avoid potential injury.
- Application of Brake to fields which have been leveled (except water leveling) within 12 months prior to application may result in serious rice injury in areas that have been cut or filled.
- Thoroughly clean the spray system with water and a suitable tank cleaner before and after each use.
- To provide adequate coverage, it is recommended that ground speed not exceed 10 mph during application.
- Avoid overlapping spray swaths as injury may occur to rotational crops.
- For best results, organic fertilizer must be incorporated into soil prior to application of Brake to the soil surface.

Application Rates

Apply this product as a postemergence residual application from the 3-leaf stage in dry-seeded rice. Refer to table below for specific rates by soil type. Application of this product to soils with high clay content (e.g. >55%) or high organic matter (e.g. >3%) may result in reduced efficacy. Minor chlorosis and stunting may occur, especially if heavy rainfall occurs during or soon after application, but plants typically outgrow these effects and develop normally.

Application Rates for Brake in Rice		
Soil Texture	Rate in Fluid Ounces per Acre	Application Timing
Sand, Loamy sand, and Sandy loam	12	Postemergence from the 3-leaf Stage
All other soil types	16	

Application Methods

Postemergence applications of this product will not control weeds that germinated prior to application. If weeds have germinated prior to application of this product, tank-mix this product with a herbicide that is effective at controlling the target weed species and compatible with the variety planted.

Postemergence Applications in Rice

This product may only be used for postemergence applications in rice. Do not apply prior to the 3-leaf stage. Do not apply more than 16 fluid ounces of this product per acre.

Moisture is necessary to activate this product in soil for residual weed control. Dry weather following applications of this product may reduce effectiveness. For improved weed control, a minimum of 0.5 inches of rainfall or irrigation is recommended following application.

Read the label affixed to the container for Brake before applying. Use of Brake according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for Brake.

New use directions appear on this supplemental label that may be different from those that appear on the container label.

© Copyright 2025 SePRO Ag, LLC

® Brake is a registered trademark of SePRO Corporation.

SePRO Ag, LLC 11550 North Meridian Street, Suite 600, Carmel, IN 46032

FPL20241219